

Steering Distillation Processes through Zeno Effect

B. Militello,¹ H. Nakazato,² and A. Messina¹

¹*INFN, MIUR and Dipartimento di Scienze Fisiche ed Astronomiche dell'Università di Palermo, Via Archirafi, 36, I-90123 Palermo, Italy*

²*Department of Physics, Waseda University, Tokyo 169-8555, Japan*

A quantum system in interaction with a repeatedly measured one undergoes a non-unitary time evolution pushing it into a subspace substantially determined by the two-system coupling. The possibility of suitably modifying such an interaction through the generalized Quantum Zeno Effect addressing the system toward an *a priori* decided target subspace is reported.

PACS numbers: 03.65.Xp, 03.65.Ta, 32.80.Pj

A distillation process, in its essence, is nothing but a systematic way of driving a system toward a state nonorthogonal to its initial condition. In some sense thus, it often represents the realization of a projection operator. In recent years, efforts have been made to realize distillation procedures since they might be exploited to prepare and to control at one's will the state of a quantum system. In the field of quantum technology[1] covering such research areas like quantum computation, quantum information or quantum teleportation it is, for instance, usually assumed that a desired state, which is necessary to start with a specific quantum manipulation, can be in principle prepared. In reality, however, it is not at all trivial and we have to specify explicitly how such a preparation is actually implemented. Various distillation processes have been proposed for this purpose so far [2].

Recently a general and new strategy of distillation specifically addressed to bipartite systems has been reported [3]. Its goal is to drive the subsystem of interest (called *Slave*) toward a final state by repeated measurement acts performed on the auxiliary subsystem (called *Master*), in interaction with the former, provided the latter part is always found in the same state. This conditional procedure, inspired by the Quantum Non-Demolition Measurements (QND) [4], shares with the latter the difficulty of singling out the suitable Master-Slave coupling allowing the selection of a *prefixed* target state. Such an intrinsic weakness poses limitations on the practical application of the method.

In this letter we propose an effective way of overcoming this structural drawback. Our scope is indeed to demonstrate the systematic possibility of controlling the distillation mechanism, furnishing conditions on the coupling to reach prefixed target subspaces. The key tool to suitably modify the two-subsystem interaction is the generalized Quantum Zeno Effect (QZE) [5, 6, 7].

Consider two interacting systems, referred to as *Master* (M) and *Slave* (S). Let $\{|\phi_k\rangle\}$ denote a basis of M , and $\{|\varphi_n\rangle\}$ indicate a basis of S . Let us denote by $\hat{U}(\tau) = e^{-i\hat{H}\tau}$ the time evolution operator ($\hbar = 1$) for time τ , $\hat{H} = \hat{H}_0$ being the compound system hamiltonian.

Perform a measurement of the Master state, and as-

sume that the result is $|\phi_0\rangle$. Let the system evolve for time τ under $\hat{H}$, then perform another measurement of the Master, and so on N times. Assume that at each step the Master system is found into the state $|\phi_0\rangle$. Under such a hypothesis the compound system is subjected to a non-unitary time evolution described by

$$\begin{aligned}\hat{W}^{(N)}(\tau) &\equiv \mathbb{N}_N \left[|\phi_0\rangle \langle \phi_0| \hat{U}(\tau) \right]^N |\phi_0\rangle \langle \phi_0| \\ &= \mathbb{N}_N |\phi_0\rangle \langle \phi_0| \left[\hat{V}(\tau) \right]^N,\end{aligned}\quad (1)$$

where

$$\hat{V}(\tau) \equiv \langle \phi_0| e^{-i\hat{H}\tau} |\phi_0\rangle \quad (2)$$

is a non-unitary operator describing the transformation that the Slave state undergoes owing to both the time evolution $\hat{U}(\tau)$ and the projection following the successful measurement act on M . The normalization constant $\mathbb{N}_N$ takes into account the probability of finding M in the state $|\phi_0\rangle$ N times.

Let us solve the eigenvalue problem related to $\hat{V}(\tau)$. Since this operator is not hermitian, generally its right and left eigenvalue problems turn out to have different solutions. Assume now that it is possible to *diagonalize* the operator $\hat{V}(\tau)$ in the standard way [8]

$$\hat{V}(\tau) = \sum_k \gamma_k \hat{P}_k, \quad (3)$$

where $\{\hat{P}_k\}$ are orthogonal projection operators satisfying the completeness relation $\sum_k \hat{P}_k = \hat{1}_S$, $\hat{1}_S$ being the Slave identity operator.

Introducing $\gamma = \max_k \{|\gamma_k|\}$, for large enough N , one finds

$$\hat{W}^{(N)}(\tau) \approx \mathbb{N}_N |\phi_0\rangle \langle \phi_0| \sum_{k:|\gamma_k|=\gamma} \gamma_k^N \hat{P}_k \quad (4)$$

If a unique eigenvalue of $\hat{V}(\tau)$ having modulus γ exists and in addition it is non-degenerate, then the distillation procedure realizes just a 'single-state selection.'

The success of such a procedure is based upon the realization of some specific conditions. First of all, observe

that what we have done is implementing a projection operator into a Slave subspace determined by the specific Master-Slave interaction and by the repeatedly detected Master state. Hence in order to reach the goal of distilling a state, such a state should be *included* in the initial state. Moreover, the evolution leading to the target state is not deterministically followed by the system, instead it is a ‘conditional’ one. In fact there are N *stochastic* intermediate steps (the measurement acts), and each one of them should be done successfully, that is, at each Master measurement step the result should be $|\phi_0\rangle$ (or some other prefixed Master State, but *always the same*).

Assume that the *orthonormal* basis $\{|\varphi_n\rangle\}$ constitutes the eigenstates of $\hat{V}(\tau)$. Accordingly, the action of the unitary operator $\hat{U}(\tau)$ on the state $|\varphi_n\rangle \otimes |\phi_0\rangle$ induces transitions to other Master states but leaves unchanged the Slave state which factorizes $|\phi_0\rangle$, i.e., $|\varphi_n\rangle \otimes |\phi_0\rangle \rightarrow \sqrt{\wp_n(\tau)} e^{i\xi_n(\tau)} |\varphi_n\rangle \otimes |\phi_0\rangle + \sum_{j \neq 0} \sum_m c_{j,m}(\tau) |\varphi_m\rangle \otimes |\phi_j\rangle$, $\wp_n(\tau)$ being the relevant survival probability. Therefore one immediately finds

$$\hat{V}(\tau) = \sum_n \sqrt{\wp_n(\tau)} e^{i\xi_n(\tau)} |\varphi_n\rangle \langle \varphi_n|. \quad (5)$$

Such a decomposition gives a precise physical meaning to the moduli of the eigenvalues of $\hat{V}(\tau)$ and shows that the distilled (preserved) Slave states (corresponding to higher eigenvalue moduli) are those undergoing a unitary evolutions (between two measurement acts) which do not induce (or induce smaller) abandon of the initial Master states $|\phi_0\rangle$ in favor of the others $|\phi_j\rangle$, $j \neq 0$. Therefore, each state $|\varphi_n\rangle$ may be thought of as a ‘channel’ of probability loss, which may be opened or closed depending on the specific features of $\hat{U}(\tau)$.

Consider now the case wherein all the transition channels are going to be closed, i.e. all transitions from $|\phi_0\rangle$ to different Master states are hindered. Such a circumstance may occur under the presence of an *external agent*, the *freezing agent*, related to a hamiltonian $\hat{H}_{12}$ responsible for an additional coupling between Master and Slave. Hence, in the lapse between two Master detections, the dynamics of the compound system is governed by $\hat{H}_\Lambda = \hat{H}_0 + \Lambda \hat{H}_{12}$, where Λ is the strength of the new coupling term. Let us denote by $\hat{U}_\Lambda(\tau) \equiv e^{-i\hat{H}_\Lambda \tau}$ the relevant evolution operator.

It is well known that the stronger the new interaction term (i.e., the bigger Λ) is, the more the dynamics is governed by $\hat{H}_{12}$ [5]. This means that the time evolution exhibits invariant subspaces related substantially to $\hat{H}_0$ when Λ approaches 0 and those related to the structure of $\hat{H}_{12}$ as Λ grows enough [5]. Depending on the specific features of $\hat{H}_0$ and $\hat{H}_{12}$, it is possible to provide an interpretation of this change of dynamical regime in terms of the *Continuous Measurement Quantum Zeno Effect* or *Zeno Dynamics* [7]. To better understand this point of view, consider the particular case wherein the hamiltonian $\hat{H}_0$ governs Slave-state-dependent oscillations be-

tween the two Master states $|\phi_0\rangle$ and $|\phi_1\rangle$, while $\hat{H}_{12}$ couples states characterized by the Master states $|\phi_1\rangle$ and $|\phi_2\rangle$. In this case the second coupling term may be thought of as an *observer* that, watching at the system following the unperturbed dynamics governed by $\hat{H}_0$, is able to freeze it into the initial Master state $|\phi_0\rangle$ [7]. The larger the ‘watching-strength’ (Λ) is, the stronger the ‘freezing’ effect is. Since $\hat{H}_{12}$ is, generally speaking, responsible for a Slave-state-dependent quantum Zeno effect, the distillation may be made ineffective/effective, depending on the corresponding Hilbert subspaces of the Slave.

The ability of controlling the ‘channel opening/closing’ is improvable when the more general case of *Hierarchically Controlled Dynamics* [5, 9] is considered, instead of the Zeno effect. In fact, the action of a subsequent coupling, let us denote it by $\hat{H}_{23}$ assuming for simplicity that it involves Master states $|\phi_2\rangle$ and $|\phi_3\rangle$ only, can hinder the hindering effect of $\hat{H}_{12}$. As a very simplified toy-model clarifying this point, let us consider, parenthetically, the following 4×4 tridiagonal matrix, expressing a coupling scheme between the states of a four-level system:

$$\Xi = \begin{pmatrix} 0 & \Omega & 0 & 0 \\ \Omega^* & 0 & \Lambda & 0 \\ 0 & \Lambda^* & 0 & \Gamma \\ 0 & 0 & \Gamma^* & 0 \end{pmatrix}. \quad (6)$$

The four-level system, whose relevant dynamics is here described, performs Rabi oscillations between the two lowest states (those coupled by Ω) when $\Lambda = \Gamma = 0$. Such a dynamical regime corresponds to the unperturbed time evolution. As Λ is made non-vanishing, the dynamics becomes more and more complicated until the condition $\Lambda \gg \Omega$ is reached. In such a situation, the dynamics of the lowest level is frozen, and this is just the continuous measurement quantum Zeno effect already recalled. Once Γ is also adjusted as a non-vanishing coupling constant, an unexpected phenomenon does happen: as Γ grows up, the effect due to the strong coupling Λ between the second and the third levels becomes weaker and weaker, up to the point, identified by the condition $\Gamma \gg \Lambda$, wherein the original Rabi oscillations are completely restored. In agreement with Refs. [5, 7], one can interpret Λ - and Γ -couplings as “continuous” measurements so that it is possible to give the following *metaphorical* statement: “a watched pot never boils” ($\Lambda \gg \Omega$, $\Gamma = 0$) but “a watched cook can freely watch a boiling pot” ($\Lambda \gg \Omega$, $\Gamma \gg \Lambda$) [5].

In passing, we mention the fact that the *hierarchical chain* of interactions in principle may be *extended* (with further rings), maintaining the same substantial features: the last ring of the chain is able to destroy the effects of the previous one depending on how the relevant coupling strengths compare to each other. For instance we could add a fifth level and a fourth coupling involving the fourth

and fifth levels. In this way, we obtain a 5×5 tridiagonal matrix. In concomitance with an increase of the fourth-coupling strength, the hindering of inhibition given by the third coupling (Γ) is hindered.

Coming back to our original problem, since the third coupling, $\hat{H}_{23}$, is Slave-state-dependent too, it can differently exert its effect subspace by subspace. Such a circumstance, of course, enlarges our ability of controlling the dynamical behavior of the compound system and hence the distillation process.

In the light of such considerations, we can express the main result of this paper as follows. A continuous measurement quantum Zeno effect, involving Master states in a Slave-state-dependent way, is able to hinder the $\hat{H}_0$ -governed unperturbed dynamics, eventually closing channels for Slave-state probability loss, which are open in the unperturbed dynamics. Hence the final distillation result may be predicted and, moreover, *a priori decided*, by exploiting the very clear mechanism of channel-closing or opening in connection with our will of *preserving* or *non-preserving* the corresponding Slave states. Moreover, we can go through a hierarchically controlled dynamics of the compound Master-Slave system, improving our ability in implementing the filter.

Let us now consider, as an example of this strategy, a three-level system (the Master) coupled to a harmonic oscillator (the Slave). Such a physical system may be realized in the context of trapped ions [10].

As is well known, a time-dependent quadrupolar electric field is able to confine a charged particle, providing an effective quadratic potential that induces a harmonic motion. When the confined particle is an ion, the complete result is a compound system possessing both fermionic and bosonic degrees of freedom, the first ones describing the internal motion of the electrons with respect to the atomic nucleus, the second ones describing the ion center-of-mass motion. In most of the experiments, only a few atomic states are really involved in the dynamics and a single vibrational mode is considered. Following Ref. [10], it is possible to realize an experimental setup which involves, in the dynamics, only the following three atomic levels, $|g\rangle := |^2S_{1/2}, F=1, m_F=1\rangle$, $|e_1\rangle := |^2S_{1/2}, F=2, m_F=2\rangle$, $|e_2\rangle := |^2S_{1/2}, F=2, m_F=1\rangle$, using a magnetic field of 1 mT to obtain the useful level splittings ($\omega(e_2) - \omega(e_1) \approx 100$ MHz, $\omega(g) - \omega(e_1) \approx 1$ GHz), and exploiting the auxiliary level $|^2P_{1/2}, F=2, m_F=2\rangle$ to realize Raman coupling schemes.

Consider now the action of two effective (i.e., implemented via Raman schemes) lasers, one tuned to the p th blue or red sideband of the atomic transition $|g\rangle \rightarrow |e_1\rangle$, and the other to the q th blue or red sideband of the atomic transition $|e_1\rangle \rightarrow |e_2\rangle$. The relevant interaction-picture hamiltonian in the *Rotating Wave Approximation*

is given by[11]

$$\hat{H}_\Lambda = \Omega [f_p(\hat{a}^\dagger \hat{a}, \eta_1) \hat{a}^p |e_1\rangle \langle g| + \text{H.c.}] + \Lambda [f_q(\hat{a}^\dagger \hat{a}, \eta_2) \hat{a}^q |e_2\rangle \langle e_1| + \text{H.c.}], \quad (7)$$

where p and q are integer numbers related to the specific choice of the laser frequencies (sidebands), Ω and Λ are coupling constants proportional to the laser intensities, η_j ($j = 1, 2$) are the relevant Lamb-Dicke parameters expressing the ratio between the vibrational ground state oscillation amplitudes and the laser wavelengths, $\hat{a}$ is the harmonic-oscillator annihilation operator (here the notation is such that $\hat{a}^{-s} \equiv \hat{a}^{\dagger s}$ for positive s), while $|g\rangle$, $|e_1\rangle$ and $|e_2\rangle$ are the three atomic states effectively involved in the dynamics. The functions f_p and f_q express nonlinear vibrational energy dependence of vibronic couplings and are such that for very small Lamb-Dicke parameters ($\eta \ll 1$), they almost approach unity, while for larger values of η they exhibit a strongly nonlinear behavior in the variable $\hat{a}^\dagger \hat{a}$, and possesses some zeroes.

Considering the unitary evolution due to $\hat{H}_\Lambda$ in (7) in the special regime $\Lambda = 0$ and $p = 0$, and repeatedly detecting the atomic state $|g\rangle$ leads to the standard quantum non-demolition measurements, by which it is possible to extract, i.e. distill, a number state [3, 4]. More in detail, denoting by $|n\rangle$ the harmonic oscillator Fock states, the effective non-unitary evolution operator acting upon the vibrational state is given by

$$\hat{V}_{\Lambda=0}(\tau) \equiv \langle g| e^{-i\hat{H}_{\Lambda=0}\tau} |g\rangle = \cos [\Omega f_0(\hat{a}^\dagger \hat{a}, \eta_1)\tau] = \sum_n \cos [\Omega f_0(n, \eta_1)\tau] |n\rangle \langle n|. \quad (8)$$

Since in this case the Fock states are eigenstates of $\hat{V}(\tau)$, we have $|\varphi_n\rangle := |n\rangle$. It is possible to choose $\tau = \tau_{\bar{n}}$ such that $\Omega f_0(\bar{n}, \eta_1)\tau_{\bar{n}} = \pi$. Taking into account that for $n \neq \bar{n}$, $f_0(n, \eta_1)$ and $f_0(\bar{n}, \eta_1)$ are incommensurable, the eigenvector $|\bar{n}\rangle \otimes |g\rangle$, and *it only*, is preserved by the distillation procedure ($|\cos [\Omega f_0(\bar{n}, \eta_1)\tau]| = 1$), while all the others are partially destroyed at each atomic state detection ($|\cos [\Omega f_0(n, \eta_1)\tau]| < 1$ for $n \neq \bar{n}$).

Consider now the effect of the freezing agent, i.e., assume $\Lambda \neq 0$. In this case the total Hamiltonian, $\hat{H}_\Lambda$, is substantially characterized by three-dimensional invariant subspaces, $\{|n\rangle \otimes |g\rangle, |n-p\rangle \otimes |e_1\rangle, |n-p-q\rangle \otimes |e_2\rangle\}$, wherein the operator may be represented as a 3×3 block of the form

$$\Xi_n = \begin{pmatrix} 0 & \Omega_n & 0 \\ \Omega_n^* & 0 & \Lambda_n \\ 0 & \Lambda_n^* & 0 \end{pmatrix}, \quad (9)$$

with $\Omega_n = \Omega f_p(n-p, \eta_1) \sqrt{n!/(n-p)!}$, $\Lambda_n = \Lambda f_q(n-p-q, \eta_2) \sqrt{(n-p)!/(n-p-q)!}$. Of course, depending on p and q , there could exist also invariant doublets and singlets. For instance, in correspondence to $n-p < 0$

there is a singlet, while if $n - p \geq 0$ and $n - p - q < 0$ there is a doublet.

It is straightforward to evaluate the non-unitary operator $\hat{V}_\Lambda \equiv \langle g | \hat{U}_\Lambda(\tau) | g \rangle$. In particular, in the case $p = 0$ (previously analyzed in absence of freezing agent), we obtain

$$\hat{V}_\Lambda(\tau) = \sum_{n=0}^{q-1} \cos[\Omega f_0(n, \eta_1)\tau] |n\rangle \langle n| + \sum_{n=q}^{\infty} \frac{|\Lambda_n|^2 + |\Omega_n|^2 \cos(\omega_n \tau)}{|\Lambda_n|^2 + |\Omega_n|^2} |n\rangle \langle n| \quad (10)$$

with $\omega_n = \sqrt{|\Lambda_n|^2 + |\Omega_n|^2}$. The operator is diagonal in the Fock basis.

It is easy to see that in correspondence to continuous measurement quantum Zeno effect ($\Lambda_n \gg \Omega_n$) in the n th subspace (the subspace the state $|n\rangle \otimes |g\rangle$ belongs to), the channel of probability loss is closed, otherwise it could be open. Therefore, the ability of controlling the dependence on n of the quantum Zeno effect incoming, plays the role of a real *handle grip* by which it is possible to open and close distillation channels. As a specific application, consider the dynamical regime characterized by $\Lambda \gg \Omega$, $p = 0$ and, in order to have $f_q(n - q, \eta_2)$ almost constant and non-vanishing for all Fock states, $\eta_2 \ll 1$. Under such hypotheses, $\Lambda_n \gg \Omega_n$, $\forall n$. If the measurement interval τ is not fine-tuned, i.e., $\tau \neq \tau_0^{(k)}, \tau_1^{(k)}, \dots, \tau_{q-1}^{(k)}$, $\forall$ integer k , where $\tau_j^{(k)}$ satisfies $\Omega f_0(j, \eta_1) \tau_j^{(k)} = k\pi$, all Fock states with an excitation number less than q are eliminated in the course of repeated measurements, hence realizing the projector $\hat{1} - \sum_{n=0,1,\dots,q-1} |n\rangle \langle n|$. That is,

$$[\hat{V}_\Lambda(\tau)]^N \xrightarrow{\text{large } N} \hat{1} - \sum_{n=0,1,\dots,q-1} |n\rangle \langle n|. \quad (11)$$

As another specific application, consider the implementation of the projector $\hat{1} - |\bar{n}\rangle \langle \bar{n}|$. In order to reach the goal, we consider the case where $\Lambda \gg \Omega$, $p = q = 0$, η_2 large enough to make the zeroes of f_0 *visible* also for not extremely high $\bar{n}$: $\Lambda_{\bar{n}} = \Lambda f_0(\bar{n}, \eta_2) = 0$, and τ is such that $\cos(\omega_{\bar{n}}\tau) \neq 1$. In this case, all channels result to be closed (because of the condition $\Lambda_n \gg \Omega_n$, for $n \neq \bar{n}$) except for that related to the $\bar{n}$ th subspace (being $\Lambda_{\bar{n}} = 0$).

In conclusion, in this paper we have addressed the general problem of how to project a quantum system in a desired *prefixed* subspace. We have indeed improved the distillation approach based upon repeated measurements on a part of a bipartite system[3] introducing the generalized Quantum Zeno Effect as a control mechanism

of the two-subsystem-interaction determining the target subspace. In other words, our main and novel result is that the presence of such a control mechanism acts upon the interaction transforming the original distillation process into a driven distillation. We conclude emphasizing that the method and the idea presented in this letter are general and then exploitable in different physical context both for fundamental and technological scopes.

This work is partly supported by the bilateral Italian-Japanese project 15C1 on “Quantum Information and Computation” of the Italian Ministry for Foreign Affairs, by a Grant for The 21st Century COE Program (Physics of Self-Organization Systems) at Waseda University and a Grant-in-Aid for Priority Areas Research (B) (No. 13135221), both from the Ministry of Education, Culture, Sports, Science and Technology, Japan, and by a Grant-in-Aid for Scientific Research (C) (No. 14540280) from the Japan Society for the Promotion of Science.

-
- [1] For reviews, see: A. Galindo and M.A. Martin-Delgado, Rev. Mod. Phys. **74**, 347 (2002); D. Bouwmeester, A. Ekert and A. Zeilinger, Eds., *The Physics of Quantum Information* (Springer, Berlin, 2000); M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
 - [2] C. H. Bennett *et al.*, Phys. Rev. Lett. **76**, 722 (1996); **78**, 2031(E) (1997); Phys. Rev. A **54**, 3824 (1996); J. I. Cirac, A. K. Ekert and C. Macchiavello, Phys. Rev. Lett. **82**, 4344 (1999). See also Refs. [1] and references therein.
 - [3] H. Nakazato, T. Takazawa and K. Yuasa, Phys. Rev. Lett. **90** 060401 (2003).
 - [4] R.L. de Matos Filho and W. Vogel, Phys. Rev. Lett. **76** 4520 (1996).
 - [5] P. Facchi and S. Pascazio, Phys. Rev. Lett. **89** 080401 (2002).
 - [6] H. Nakazato, M. Namiki and S. Pascazio, Int. J. Mod. Phys. B **10**, 247 (1996); D. Home and M. A. B. Whitaker, Ann. Phys. **258**, 237 (1997).
 - [7] P. Facchi and S. Pascazio, *Quantum Zeno and Inverse quantum Zeno effects*, in Progress in Optics **42**, edited by E. Wolf (Elsevier, Amsterdam, 2001), p. 147.
 - [8] The most general case, in which the standard form (3) is no longer valid, owing to the presence of degenerate eigenvalues, is found in: H. Nakazato, M. Unoki and K. Yuasa, “Preparation and entanglement purification of qubits through Zeno-like measurements,” WU-HEP-04-01 (2004), accepted for publication in Phys. Rev. A, quant-ph/0402182.
 - [9] B. Militello, A. Messina and A. Napoli, Fortschr. Phys. **49**, 1041 (2001).
 - [10] S. Maniscalco and A. Messina, Fortschr. Phys. **49**, 1027 (2001).
 - [11] B. Militello, A. Messina and A. Napoli, Phys. Lett. A **286**, 239 (2001).